



UNIVERSITI PUTRA MALAYSIA

**NUTRIENT COMPOSITION AND HYPOCHOLESTEROLEMIC EFFECT
OF BLACK SEEDS (*NIGELLA SATIVA* L.) AND ITS OIL**

GHANYA NAJI AL-NAQEEB

FPSK(M) 2005 16

**NUTRIENT COMPOSITION AND HYPOCHOLESTEROLEMIC EFFECT OF
BLACK SEEDS (*NIGELLA SATIVA L.*) AND ITS OIL**

By

GHANYA NAJI AL-NAQEEB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

May 2005



*To may parent
To Al-Naoeep family
To all people from Saber*

*To my country, Republic of Yemen
To those who are really very hard working in any where*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**NUTRIENT COMPOSITION AND HYPOCHOLESTEROLEMIC EFFECT OF
“BLACK SEEDS” (*NIGELLA SATIVA* L.) AND ITS OIL**

By

GHANYA NAJI AL-NAQEEB

May 2005

Chairman: Associate Professor Maznah Ismail, Ph.D.

Faculty : Medicine and Health Sciences

Nigella sativa Linn. (Ranunculacea) popularly known as the black seed or Habbatus Sauda is a herb that has traditionally been used for centuries in many parts of the world. It has gained popularity due to its potential health benefits. This study was carried out to investigate the nutrient composition and *in vivo* hypocholesterolemic effect of the black seeds cultivated in Yemen and its oil. The nutrient composition of three different samples of *Nigella sativa* seeds from three different regions in Yemen, namely Marib, Sadah and Taiz were studied. Proximate analysis, total dietary fiber (TDF), insoluble dietary fiber (IDF) and soluble dietary fiber (SDF) were determined by the Association of Official Analytical Chemists methods (AOAC). Mineral analysis was carried out using Atomic Absorption spectrophotometer (AAS). Fatty acid methyl esters composition was determined using Gas Chromatography (GC), vitamin E (α -tocopherol) using HPLC and antioxidant activity by ferric thiocyanate (FTC) and thiobarbituric acid (TBA) methods. Results show that the three samples of *Nigella sativa* seeds have high content of fat which were 38.4 ± 2.1 , 37.7 ± 0.1 , and $36.8 \pm 1.4\%$ in Sadah, Marib, and Taiz, respectively. In addition, *Nigella sativa* seeds have a high content of TDF (36.88 ± 1.44 , 26.50 ± 1.05 , $30.40 \pm 1.06\%$ for Marib, Sadah and Taiz

samples, respectively), IDF (27.10 ± 0.55 , 20.56 ± 1.16 , 22.40 ± 1.40 % for Marib, Sadah and Taiz samples, respectively) and SDF (8.90 ± 1.17 , 6.50 ± 0.60 , 8.13 ± 0.71 % for Marib, Sadah and Taiz samples, respectively).

Nigella sativa seeds are also rich in calcium, magnesium, potassium, phosphorus and iron. The major fatty acids in *Nigella sativa* oil extracts from n-hexane were oleic and linolenic which exist as unsaturated fatty acids. Palmitic, stearic and myristic acids were the main saturated fatty acids in all three samples. Oil extracted from the seeds showed higher antioxidant activity compared to α -tocopherol, and was rich in vitamin E (120 ± 0.15 , 170 ± 0.40 and 290 ± 1.5 mg/100g for Marib, Sadah and Taiz samples, respectively).

The effect of *Nigella sativa* seed powder and its oil on hypercholesterolemia-induced rabbits was also studied for 8 weeks. A significant reduction ($p < 0.05$) of total cholesterol (TC) and low density lipoproteins (LDL) in *Nigella sativa* powder (NSP) and *Nigella sativa* oil (NSO) groups were observed at weeks 2, 4, 6 and 8. Treatment of rabbits with NSP and NSO showed a significant increase ($p > 0.05$) in plasma HDL levels at weeks 4, 6 and 8 of treatment. A significant decreased ($p < 0.05$) of malondialdehyde (MDA) concentration in erythrocyte and plasma was observed in NSP and NSO groups. There was a significant increase of total plasma antioxidant status (TAS) ($p > 0.05$). NSP and NSO groups also showed significant reduction of serum alanine aminotransferase (ALT) and creatinine concentrations ($p < 0.05$). Plaque formation was significantly inhibited in the aorta, and the tunica intima to tunica media ratio was significantly decreased ($p < 0.05$).

In conclusion, this study indicates that the seed of *Nigella sativa* cultivated in Yemen and its oil contain several nutrients and antioxidants with potential health values such as hypocholesterolemic effect, antioxidant effect and anti atherogenic effect.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

KOMPOSISI NUTRIEN DAN KESAN HIPOKOLESTEROLEMIK “BIJAN HITAM” (*NIGELLA SATIVA* L.) DAN MINYAKNYA

Oleh

GHANYA AL-NAQEEB

May 2005

Pengerusi: Profesor Madya Maznah Ismail, Ph.D.

Fakulti : Perubatan dan Sains Kesihatan

Nigella sativa L. (Ranunculacea) yang biasanya dikenali sebagai bijan hitam atau Habbatus Sauda merupakan sejenis herba yang digunakan secara meluas sejak berkurun lamanya di kebanyakan negara di dunia disebabkan oleh potensi kesihatannya. Walau bagaimanapun, lebih banyak maklumat saintifik diperlukan bagi menyokong potensi kesihatan ini. Kajian ini dijalankan untuk mengkaji komposisi nutrien serta kesan hipokolestrolemik bijan hitam yang ditanam di Yemen dan minyaknya secara *in vivo*. Komposisi nutrien telah dianalisis daripada tiga sampel biji *Nigella sativa* dari tiga kawasan iaitu Marib, Sadah dan Taiz. Analisis proksimat, jumlah fiber diet (TDF), fiber diet tidak larut (IDF) dan fiber diet larut telah ditentukan melalui kaedah AOAC. Analisis mineral (besi, kalsium, magnesium, kalium, natrium, fosforus dan zink) telah dijalankan dengan menggunakan sistem nyalaan, Spektrofotometer Serapan Atom (AAS). Kandungan asid lemak metil ester telah ditentukan melalui kaedah kromatografi gas (GC-MS), kandungan vitamin E (α -tokoferol) menggunakan kromatografi cecair prestasi tinggi (HPLC), dan kandungan antioksidan melalui kaedah ferik tiosianat (FTC) dan asid tiobarbiturik (TBA).

Keputusan menunjukkan ketiga-tiga sampel biji *Nigella sativa* mengandungi kandungan minyak yang tinggi, iaitu 37.7 ± 1.5 , 38.4 ± 2.1 dan $36.8 \pm 1.4\%$, masing-masing dalam sampel Marib, Sadah dan Taiz. *Nigella sativa* juga mempunyai kandungan TDF yang tinggi, iaitu 36.88 ± 1.44 , 26.50 ± 1.05 , $30.40 \pm 1.06\%$, IDF, iaitu 27.10 ± 0.55 , 20.56 ± 1.16 , $22.40 \pm 1.40 \%$, dan SDF, iaitu 8.90 ± 1.17 , 6.50 ± 0.60 , $8.13 \pm 0.71\%$ masing-masing untuk sampel Marib, Sadah dan Taiz.

Nigella sativa kaya dengan kalsium, magnesium, kalium, fosforus dan zat besi. Komposisi asid lemak metilester bagi minyak yang diekstrak menggunakan *n*-heksana ialah oleik dan linoleik yang hadir sebagai asid lemak tak tepu yang utama. Asid palmitik, stearik dan miristik merupakan asid lemak tepu yang utama dalam ketiga-tiga sampel dari Marib, Sadah dan Taiz. Minyak daripada biji *Nigella sativa* telah menunjukkan aktiviti antioksidan yang lebih tinggi berbanding α -tokoferol rujukan, dan didapati kaya dengan vitamin E (120 ± 0.15 , 170 ± 0.40 dan 290 ± 1.5 mg/100g masing-masing bagi sampel Marib, Sadah dan Taiz).

Kesan serbuk biji dan minyak bijan hitam kepada arnab yang diaruh hiperkolestrolemik telah dikaji selama 8 minggu. Penurunan yang signifikan ($p < 0.05$) bagi jumlah kolesterol (TC), lipoprotein ketumpatan rendah (LDL) dalam kumpulan diberi serbuk *Nigella sativa* (NSP) dan kumpulan yang diberikan minyak *Nigella sativa* (NSO) berlaku pada minggu rawatan ke 2, 4, 6 dan 8. Arnab yang dirawat dengan NSP dan NSO menunjukkan peningkatan yang signifikan ($p < 0.05$) pada tahap lipoprotein berketumpatan tinggi (HDL) dalam plasma pada minggu ke 4, 6 dan 8. Tiada perubahan yang signifikan didapati bagi paras trigliserida (TG) sepanjang masa rawatan. Kedua-

dua kumpulan NSP dan NSO menunjukkan penurunan yang signifikan ($p < 0.05$) bagi kepekatan malondialdehid (MDA) dalam eritrosit dan plasma. Jumlah status antioksidan plasma (TAS) juga telah meningkat secara signifikan ($p < 0.05$). NSP dan NSO pula menunjukkan penurunan yang signifikan bagi kepekatan alanin aminotransferase (ALT) dan kreatinin dalam serum ($p < 0.05$).

Pembentukan plak juga telah direncat dan nisbah tunika intima kepada tunika media pada lapisan dinding aorta telah menurun secara signifikan ($p < 0.05$). Kesimpulannya, kajian ini telah menunjukkan bahawa serbuk biji *Nigella sativa* yang ditanam di Yemen dan minyaknya mengandungi pelbagai nutrien dan antioksidan yang dapat memberikan kesan hipokolestolemik, antioksidan dan anti-aterogenik.

ACKNOWLEDGEMENTS

I owe first and foremost my profound gratitude to almighty ALLAH, the source of all inspiration and help, and without whose assistance, this study would not have come into existence.

Deep obligation and indebtedness and most sincere gratitude are offered to my supervisor Associate Professor Dr. MAZNAH ISMAIL for her continuous guidance during all stages of my research work and for her willingness to help, listen and assist in every way, in the midst of her heavy responsibilities. I also record my sincere appreciation to the committee members: Dr. NORHAIZAN MOHD. ESA and Dr. ZULKHAIRI HJ AMOM, for their guidance, valuable advice and precious evaluation and valuable comments.

I sincerely appreciate and acknowledge staff, laboratory assistants and graduate students at Department of Nutrition and Health Sciences for their help and cooperation. Finally I am deeply obligated and thankful to my father, my mother, my sisters, brothers, all my friends and Al – Barra for their great help, patience, support and their encouragement over the course of this thesis.

I certify that an Examination Committee met on 30th May 2005 to conduct the final examination of Ghanya Naji Al-Naqeeb on her Master of Science thesis entitled “Nutrient Composition and Hypcholesterolemic Effect of *Nigella sativa* L.” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Zaitun Yassin, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Rokiah Mohd Yusof, PhD


Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Azizah Abdul Hamid, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Suria Abd Rahman, PhD

Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **21 JUL 2005**

This thesis submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Sciences. The members of the Supervisory Committee are as followed:

MAZNAH ISMAIL, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

NORHAIZAN MOHD. ESA, Ph.D.

Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

ZULKHAIRI HJ AMOM, Ph.D.

Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)




AINI IDERIS, Ph.D.

Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **11 AUG 2005**

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



GHANYA NAJI AL-NAQEEP

Date: 18/7/2005

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	i
ACKNOWLEDGEMENTS	v
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xxi
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xviii
CHAPTER	xx
 1 INTRODUCTION	 1
 2 LITERATURE REVIEW	 6
2.1 Cholesterol	6
2.2 Lipoproteins	7
2.2.1 Chylomicrons	7
2.2.2 Very Low Density Lipoprotein (VLDL)	8
2.2.3 Low Density Lipoprotein (LDL)	8
2.2.4 High Density Lipoprotein (HDL)	9
2.3 Bile Acids	10
2.4 Atherosclerosis	11
2.4.1 Hypercholesterolemia And The Risk of Atherosclerosis	11
2.4.2 Atherosclerosis Formation	13
2.5 Lipid Peroxidation	14
2.5.1 Hypercholesterolemia And Lipid Peroxidation	15
2.6 Hypercholesterolemia And Antioxidants	17
2.7 Hypercholesterolemia And Toxicity	18
2.7.1 Alanine Aminotransferase (ALT)	18
2.7.2 Gamma –Glutamyltranspeptidase	19
2.7.3 Creatinine	19
2.7.4 Urea	20
2.8 Hydroxy-3-methylglutaryl Coenzyme A (HMG-CoA reductase)	20
2.8.1 Side Effects of Statin	22
2.9 Black Cumin Seed (<i>Nigella Sativa</i> L)	23
2.9.1 Physical And Chemical Properties of <i>Nigella</i> seeds	25
2.9.2 Nutrient Composition of <i>Nigella Sativa</i> Seeds	25
2.9.3 Vitamins Composition of <i>Nigella Sativa</i> Seeds	25
2.9.4 Amino Acid And Fatty Acid Content of <i>Nigella Sativa</i> Seeds	27
2.10 <i>Nigella Sativa</i> Oil	27
2.10.1 Fatty Acid Composition	28
2.10.2 Volatile Oil	29
2.11 Health Benefits of <i>Nigella Sativa</i> Seeds	30
2.11.1 Hypocholesterolemic Effect of <i>Nigella Sativa</i> Seeds	31



2.11.2	Effect of <i>Nigella Sativa</i> on Lipid Peroxidation	32
2.11.3	Antioxidant Effect of <i>Nigella Sativa</i> And Its Oil.	33
2.11.4	Effect of <i>Nigella Sativa</i> On Liver And Kidney Diseases	33
3	MATERIALS AND METHODS	35
3.1	Materials	35
3.1.1	<i>Nigella Sativa</i> Seeds Collection And Storage	35
3.2	Methods	35
3.2.1.	Chemical Analysis	35
3.2.2.	Experimental Designe	36
3.2.1.1	Moisture Content	38
3.2.1.2	Ash Content (Dry Ashing)	38
3.2.1.3	Crude Fat	38
3.2.1.4	Crude Protein	39
3.2.1.5	Total Dietary Fiber Analysis	39
3.2.1.6	Mineral Analysis	41
3.3	<i>Nigella Sativa</i> Seed Oil	41
3.3.1	Crude Oil Extraction	41
3.3.2	Fatty Acid Composition of The Oil Extracted	42
3.3.2.1	Gas Chromatography Condition	42
3.3.3	Total Antioxidant Activity of <i>Nigella Sativa</i> Seed Oil	43
3.3.3.1	Ferric Thiocyanate (FTC) Method	43
3.3.3.2	Thiobarbituric Acid (TBA) Method	44
3.3.4	Determination of Vitamin E of <i>Nigella Sativa</i> Oil Using HPLC	45
3.4	<i>In Vivo</i> Study	46
3.4.1	Animal Groups	46
3.4.2	Diet Ingredients and Preparation	46
3.4.3	Experimental Design	48
3.4.4	Food Intake and Body Weight	49
3.4.5	Blood Collection	49
3.4.4.1	Erythrocyte Preparation	49
3.4.5	Determination of Malondialdehyde (MDA)	50
3.4.5.1	Erythrocyte Malondialdehyde Level	50
3.4.5.2	Plasma Malondialdehyde Level	51
3.4.6	Lipid Profile Analysis	51
3.4.6.1	Analysis of Total Cholesterol (TC)	51
3.4.6.2	Analysis of High Density Lipoprotein (HDL)	52
3.4.6.3	Analysis of Low Density Lipoprotein (LDL)	52
3.4.6.4	Analysis of Triglycerides (TG)	53
3.4.7	Total Antioxidant Status (TAS) Assay	54
3.4.8	Toxicity Study	54
3.4.8.1	Alanine Transaminase (ALT) Assay	55
3.4.8.2	Gamma –Glutamyltranspeptidase (GGT) Assay	55
3.4.8.3	Creatinine Assay	56
3.4.8.4	Urea Assay	56
3.4.9	Histology Study	57
3.4.9.1	Atheroma Plaques	57
3.4.9.2	Light Microscopy Analysis	57



3.4.9.2.1	Tissue Blocking	58
3.4.9.2.2	Tissue Sectioning	58
3.4.9.2.3	Tissues Staining	59
3.4.9.3	Quantitative Analysis of Histological Data	60
3.5	Statistical Analysis	60
4	RESULTS AND DISCUSSION	61
4.1	Nutrient Composition of Three Different Samples of <i>Nigella Sativa</i>	61
4.1.1	Proximate Analysis of <i>Nigella Sativa</i> Seeds	61
4.1.2	Fiber Content	64
4.1.3	Minerals Content	66
4.2	<i>Nigella Sativa</i> Oil	69
4.2.1	Fatty Acid Composition	70
4.2.2	Total Antioxidants Activity of <i>Nigella sativa</i> Seed Oil	74
4.2.2.1	Ferric Thiocyanate Method	74
4.2.2.2	Thiobarbituric Acid Method	76
4.2.3	α -tocopherol Content of <i>Nigella Sativa</i> Seed Oil	80
4.3	Animal Study	81
4.3.1	Body Weight	81
4.3.2	Liver And Kidney Weight of Experimental Rabbits	83
4.3.3	Lipid Profile	84
4.3.3.1	Total Cholesterol levels	84
4.3.3.2	Low Density Lipoprotein Level	85
4.3.3.3	High Density Lipoproteins Level	88
4.3.3.4	Triglycerides Level	88
4.3.4	Lipid Peroxidation	93
4.3.4.1	Malondialdehyde Level In Erythrocyte	94
4.3.4.2	Malondialdehyde Level In Plasma	95
4.3.5	Total Antioxidant Status (TAS)	99
4.3.6	Toxicity Study	100
4.3.6.1	Alanine Aminotransferase (ALT)	100
4.3.6.2	Gamma-Glutamyl Transferase (GGT)	103
4.3.6.3	Creatinine Level	104
4.3.6.4	Urea Concentration In Serum	107
4.3.7	Histology Study	110
4.3.7.1	Assessment of Atherosclerotic Plaques	110
4.3.7.2	Quantitative Analysis of Histological Data.	113
4.3.7.2.1	Intima and Media Thickness	113
4.3.7.2.2	Intima: Media Ratio	117
5	CONCLUSION	126
6	RECOMMENDATIONS	129

REFERENCES	130
APPENDICES	141
BIODATA OF THE AUTHOR	148



LIST OF TABLES

Table		Page
2.1	Some characteristics of <i>Nigella sativa</i> seeds, as reported in the literature	26
2.2	Mineral contents of <i>Nigella sativa</i> seeds as reported in the literature	26
2.3	Vitamin composition of <i>Nigella sativa</i> seeds as reported in the literature	27
2.4	Amino acid content of <i>Nigella sativa</i> seeds as reported in the literature	29
2.5	Quantification of quinones in oil from <i>Nigella sativa</i> seed	30
3.1	The ingredient of the diet prepared	47
3.2	Tissue Dehydration in a tissue processor machine	58
3.3	Colouration with Hematoxyline and Eosin (H&E)	59
4.1	Nutrient composition of three different samples of <i>Nigella sativa</i> seeds cultivated in Yemen.	62
4.2	Proximate analysis of <i>Nigella sativa</i> seeds in comparison with selected nuts and oil seeds.	64
4.3	Fiber content of defatted <i>Nigella sativa</i> seeds	65
4.4	Mineral content of three different samples of <i>Nigella sativa</i> seeds cultivated in Yemen and some results reported in the literature	67
4.5	Oil yield from three different samples of <i>Nigella sativa</i> seeds	70
4.6	Fatty acid composition of three different samples of <i>Nigella sativa</i> oil	72
4.7	Percentage of antioxidant activity of three different samples of <i>Nigella sativa</i> seed oil extracted using three different solvents. (FTC and TBA method).	75
4.8	α -Tocopherol content of three different samples of <i>Nigella sativa</i> seed oil	81
4.9	Livers and kidneys weights	83

		87
4.11	Changes in plasma LDL levels during the experimental period	
4.12	Changes in plasma triglycerides Level of experimental rabbits	90
4.13	Changes of MDA levels in erythrocytes of the rabbits over experiment period	96
4.14	Changes of plasma MDA levels of the rabbits over the experiment period	97
4.15	Total antioxidant status of plasma after induction with 0.5% cholesterol for 3 weeks and after 8 weeks of treatment	100
4.16	Changes in serum creatinine levels t of experimental rabbits	106
4.17	Changes in serum urea levels of rabbits during the experimental period	109
4.18	Percentage of lesion area in experimental rabbits	111
4.19	Maximum thickness of the intima, media and intima to media ratio in experimental rabbits	117

LIST OF FIGURES

Figure		Page
2.1	Structure of low-density lipoprotein	9
2.2	Changes in the arterial intima during atherogenesis	16
3.1	Samples sources (Marib, Sadah and Taiz) in Yemen	36
3.2	Flow-chart for the research design	37
3.3	Animals group	48
4.1	Absorbance values of <i>Nigella sativa</i> seed oil samples from Marib using FTC method.	78
4.2	Absorbance values of <i>Nigella sativa</i> seed oil samples from Sadah using TBA method.	79
4.3	Changes in body weights of experimental rabbits	82
4.6	Changes in plasma HDL levels of the experimental rabbits	89
4.5	Serum Alanine Aminotransferase concentration	101
4.6	Serum Gamma-Glutamyl Transferase concentration	105
4.7	Intimal surface of the aorta from NC group	112
4.8	Intimal surface of the aortas from PC group showing Sudan IV stainable lipid deposit. Lipid deposits are stained brick red.	112
4.9	Intimal surface of the aortas from NSP group showing Sudan IV stainable lipid deposit. Lipid deposits are stained brick red.	113
4.10	Intimal surface of the aortas from NSO group showing Sudan IV stainable lipid deposit. Lipid deposits are stained brick red.	113
4.11	Intimal surface of the aortas from ST group showing Sudan IV stainable lipid deposit. Lipid deposits are stained brick red.	114
4.12	Hematoxylin and eosin stained of the intimal thickening of aorta for negative control group and positive control (x10)	117



4.13	Hematoxylin and eosin stained of the intimal thickening of aorta for <i>Nigella sativa</i> in powder form group and <i>Nigella sativa</i> oi	118
4.14	Hematoxylin and eosin stained of the intimal thickening of aorta for simvastatin group (x10) and positive control group (x20)	119
4.15	Hematoxylin and eosin stained of the intimal thickening of aorta for negative control and positive control group (x40)	120
4.16	Hematoxylin and eosin stained of the intimal thickening of aort for <i>Nigella sativa</i> in powder form and <i>Nigella sativa</i> oil (x40).	121
4.17	Hematoxylin and eosin stained of the intimal thickening of aorta from simvastatin group.	122
4.18	Hematoxylin and eosin stained of the intimal thickening of aort for the positive control group (x100).	123
4.19	Effect of <i>Nigella sativa</i> and simvastatin treatment on the intima and media thickness of aorta.	124

LIST OF ABBREVIATIONS

Apo	Apolipoprotein
ALT	Alanine Aminotransferase
ALP	Alkaline Phosphatase
ANOVA	One Way ANOVA
CHD	Coronary heart disease
CCL ₄	Carbon tetrachloride
FTC	Ferric Thiocyanate
GC	Gas Chromatography
GGT	Gamma Glutamyltranspeptidase
HDL	High density lipoprotein
HMG-CoA	3-hydroxy-3-methylglutaryl Coenzyme - A
IDF	Insoluble Dietary Fibre
IDL	Intermediate Density Lipoprotein
LCAT	Lecithin: Cholesterol Acyltransferase
LDL	Low Density Lipoprotein
LDH	Lactate Dehydrogenase
MDA	Malondialdehyde
min	Minute
NC	Negative control
PKC	Palm kernel cake
NSO	<i>Nigella Sativa</i> Seed Oil
SFA	Saturated Fatty Acid
SDF	Soluble Dietary Fibre
ST	Simvastatin

TAS	Total Antioxidant Status
TBA	Thiobarbituric Acid
TBARS	Thiobarbituric Acid Reactive Substances
TDF	Total Dietary Fiber
VLDL	Very Low Density Lipoprotein
USFA	Unsaturated Fatty Acid
Wk	Week



CHAPTER 1

INTRODUCTION

Cholesterol is an essential constituent of all animal cells, without it the living cells would not function properly and the organism would die (Saraswathi *et al.*, 1997). It has been documented that high plasma cholesterol level (hypercholesterolemia) is one of the most important coronary risk factors. Cardiac morbidity and mortality are directly related to serum cholesterol levels (Chen *et al.*, 1999). Many investigations indicate that elevated serum cholesterol levels may modify the biochemical properties of blood components and arterial intima, thus enhancing the development of atherosclerosis (Wallidus *et al.*, 1993).

It was reported that hypercholesterolemia induces a free radical mediated lipoprotein peroxidation and causes oxidative stress. This stress results from the imbalance between the production of free radicals and the effectiveness of the antioxidant defense system. The activity of free radicals is countered by a system of antioxidant defenses, of which vitamin E is the major chain breaking lipophilic antioxidant in tissues and plasma. In addition to vitamin E, tissues and erythrocyte enzymes contribute to the cell antioxidant defense mechanism. These include superoxide dismutase (SOD), catalase and glutathione peroxidase (GSH-Px), which detoxify H_2O_2 and convert lipid hydroperoxides to nontoxic alcohol. Disorders in these erythrocyte enzyme activities have been reported in subjects with cardiovascular disease (Halliwell *et al.*, 1996).

Reduction of cholesterol level was the main target in preventing the development of coronary heart disease (CHD). Health care plans have also included cholesterol reduction and dietary changes as a way to protect against CHD (Abajo *et al.*, 1993).

There is great interest in choosing the best diet to maintain desirable serum lipid levels, which may protect against heart disease. High levels of serum total cholesterol, low-density lipoprotein cholesterol (LDL), and triacylglycerol levels have been associated with increased risk of heart disease, while high levels of high-density lipoprotein cholesterol (HDL) have been associated with decreased risk of heart disease (Nicolosi *et al.*, 2001). Thus, a dietary pattern that most effectively lowers TC, LDL, and triacylglycerol levels, while maintaining or increasing HDL would be desirable. However, the ideal dietary intervention to improve serum lipids and reduce heart disease risk remains controversial (Nicolosi *et al.*, 2001).

It has been shown that increased consumption of omega-3 polyunsaturated fatty acids improves endothelium-dependent relaxation and protects against the development of atherosclerotic cardiovascular diseases (Prasad *et al.*, 1997; Nicolosi *et al.*, 2001). Studies have shown that vitamin E administered to hypercholesterolemic rabbits significantly reduced the plasma LDL and vessel wall oxidation, a potential beneficial action of vitamin E in hypercholesterolemic patients (Jorge *et al.*, 1998). Consumption of products containing dietary fiber has increased due to the health benefits that they offer (Khaw *et al.*, 1987). Insoluble as well as soluble dietary fibers have been associated with cholesterol lowering properties, which can be effective in lowering the risk of cardiovascular disease. Recent findings have demonstrated that the insoluble fiber derived from some fruits and